

CIRCADIAN CLOCK-ASSOCIATED 1 and LATE ELONGATED HYPOCOTYL regulate expression of the C-REPEAT BINDING FACTOR (CBF) pathway in *Arabidopsis*

Malia A. Dong^{a,b}, Eva M. Farré^b, and Michael F. Thomashow^{a,c,1}

^aMichigan State University-Department of Energy Plant Research Laboratory, ^bDepartment of Plant Biology, and ^cDepartment of Crop and Soil Sciences, Michigan State University, East Lansing, MI 48824

Contributed by Michael F. Thomashow, March 9, 2011 (sent for review February 10, 2011)

The C-REPEAT BINDING FACTOR (CBF) cold-response pathway has a prominent role in cold acclimation, the process whereby certain plants increase tolerance to freezing in response to low non-freezing temperatures. In *Arabidopsis*, the CBF pathway is characterized by rapid induction of the C-REPEAT BINDING FACTOR 1 (CBF1), CBF2, and CBF3 genes, which encode transcriptional activators, followed by induction of the CBF-targeted genes known as the “CBF regulon.” Expression of the CBF regulon results in an increase in freezing tolerance. Previous studies established that CBF1, CBF2, and CBF3 are subject to circadian regulation and that their cold induction is gated by the circadian clock. Here we present the results of genetic analysis and ChIP experiments indicating that both these forms of regulation involve direct positive action of two transcription factors that are core components of the clock, i.e., CIRCADIAN CLOCK-ASSOCIATED 1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY). In plants carrying the *cca1-11/lhy-21* double mutation, cold induction of CBF1, CBF2, and CBF3 was greatly impaired, and circadian regulation of CBF1 and CBF3 was essentially eliminated; circadian regulation of CBF2 continued, although with significantly reduced amplitude. Circadian regulation and cold induction of three CBF regulon genes, i.e., COLD-REGULATED GENE15A (COR15A), COR47, and COR78, also were greatly diminished in plants carrying the *cca1-11/lhy-21* double mutation. Furthermore, the *cca1-11/lhy-21* double mutation resulted in impaired freezing tolerance in both nonacclimated and cold-acclimated plants. These results indicate that CCA1/LHY-mediated output from the circadian clock contributes to plant cold tolerance through regulation of the CBF cold-response pathway.

A general feature of plants from temperate environments is that they increase in freezing tolerance in response to low non-freezing temperatures, a process called “cold acclimation” (1, 2). It now is well established that cold acclimation involves extensive changes in gene expression (3–6). The best understood cold-regulatory pathway is the CBF pathway. This pathway, which is widely conserved in plants (7), is best characterized in *Arabidopsis* (8, 9). When *Arabidopsis* plants are transferred from warm to cold temperature, C-REPEAT BINDING FACTOR 1 (CBF1), 2 (CBF2), and 3 (CBF3)—also known as DROUGHT RESPONSE ELEMENT BINDING FACTOR 1B (DREB1B), 1C (DREB1C), and 1A (DREB1A), respectively—are induced rapidly. These genes, which are linked physically in tandem array, encode transcription factors that are members of the AP2/ERF family of DNA-binding proteins (10). The CBF proteins bind to the CRT/DRE regulatory element present in the promoters of about 100 cold-regulated (COR) genes, known as the “CBF regulon,” and induce their expression (4, 6, 11). Constitutive overexpression of CBF1, CBF2, and CBF3 at warm temperature results in constitutive expression of the CBF regulon and an increase in freezing tolerance (12–14). The mechanisms whereby expression of the CBF regulon promotes freezing tolerance are not completely understood but involve the synthesis of low molecular weight cryoprotectants such as sucrose and raffinose and proteins that have cryoprotective properties (1, 2).

Given their importance in cold acclimation, efforts have been directed at understanding the mechanisms involved in cold-induction of CBF1, CBF2, and CBF3. To date, two positive regulators have been identified: INDUCER OF CBF EXPRESSION 1 (ICE1), a Myc family transcription factor that positively regulates CBF3 (15), and CALMODULIN-BINDING TRANSCRIPTION ACTIVATOR 3 (CAMTA3), a CAMTA family transcription factor that positively regulates CBF1 and CBF2 (16). The ICE1 and CAMTA3 genes are transcribed at warm temperature, indicating that their activities involve posttranscriptional regulatory mechanisms that are responsive to low temperature (16–18).

Another factor that affects the expression of CBF1, CBF2, and CBF3 is the circadian clock (19–22). At warm temperature, the transcript levels for CBF1, CBF2, and CBF3 oscillate with a peak at about 8 h after dawn (zeitgeber time 8; ZT8) and a trough at about ZT20. Moreover, cold-induction of CBF1, CBF2, and CBF3 is “gated” by the clock (22); if plants are exposed to low temperature at ZT4, the increase in CBF1, CBF2, and CBF3 transcript levels is much greater than if plants are exposed to low temperature at ZT16. These results indicate that cold induction of CBF1, CBF2 and CBF3 involves the integration of low-temperature and clock-regulatory pathways.

The circadian clock of *Arabidopsis* consists of multiple interlocking regulatory feedback loops (23, 24). Key components of the core feedback loop are CIRCADIAN CLOCK-ASSOCIATED 1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY), Myb transcription factors that have partially overlapping functions (25–28), and TIMING OF CAB 1 (TOC1), a PSEUDO RESPONSE REGULATOR (PRR) protein (29). Expression of CCA1 and LHY peaks just after dawn, whereas the expression of TOC1 peaks in the early evening. CCA1 and LHY bind to the Evening Element (EE) (19) present in the promoter of TOC1 and repress its transcription (30). TOC1 is necessary for the induction of both CCA1 and LHY (28). TOC1 is known to inhibit the repression of CCA1 by the TCP transcription factor CCA1 HIKING EXPEDITION 1 (CHE1) (31), but the means by which LHY expression is activated by TOC1 remains unknown. CCA1 and LHY also regulate expression of PSEUDO RESPONSE REGULATORS 7 (PRR7) and 9 (PRR9) (32, 33), two components of the morning regulatory loop. CCA1 and LHY bind to the promoters of these two genes to induce their expression, and the PRR7 and PRR9 proteins then negatively regulate CCA1 and LHY (32, 33).

Author contributions: M.A.D., E.M.F., and M.F.T. designed research; M.A.D. performed research; M.A.D., E.M.F., and M.F.T. analyzed data; and M.A.D., E.M.F., and M.F.T. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: thomash6@msu.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1103741108/-DCSupplemental.




Fig. 1. Effects of the *cca1-11/lhy-21* double mutation on circadian regulation of *CBF1*, *CBF2*, *CBF3*, and CBF-targeted genes *COR15A*, *COR47*, and *COR78*. Wild-type Wassilewskija-2 (WS) and *cca1-11/lhy-21* double-mutant plants were grown at 22 °C under a 12-h photoperiod to the four-leaf stage and then were transferred to constant light at zeitgeber time 0 (ZT0) (subjective day and night are indicated by white and gray bars, respectively). Plants were harvested every 2 h, and the transcript levels for the indicated genes were determined by qRT-PCR. Gene expression was normalized to *ISOPENTENYL PYROPHOSPHATE (IPP2)* for each sample. Gene expression is relative to one wild-type sample set to a value of 1 for each biological replicate. Values are averages from three independent biological experiments. Error bars indicate SEM.

It was reported recently that *Arabidopsis* plants carrying the *prr5/prr7/prr9* triple mutation constitutively express *CBF1*, *CBF2*, and *CBF3* at high levels and display constitutively high levels of freezing tolerance (34). Thus, it was proposed that *PRR5*, *PRR7*, and *PRR9* might act as direct negative regulators of *CBF1*, *CBF2*, and *CBF3* (34). Here we present results indicating that the clock also provides positive regulation of the CBF cold-response pathway and enhances freezing tolerance through action of the core clock components CCA1 and LHY.

Results

CCA1 and LHY Have a Direct Role in Circadian Regulation of *CBF1*, *CBF2*, and *CBF3*. Consistent with previous reports (19–21), we found that *CBF1*, *CBF2*, and *CBF3* are subject to circadian regulation (Fig. 1). Transcript levels for *CBF1*, *CBF2*, and *CBF3* oscillated with a peak occurring at about ZT8 followed by a second peak about 24 h later. The oscillation patterns for the three *CBF* genes were similar, although we observed one consistent difference; whereas the ZT8 peak for *CBF1* was lower than the

second peak, the ZT8 peaks for *CBF2* and *CBF3* were higher than the second peak. Thus, the transition from dark to light may have a specific effect on the regulation of *CBF1*.

Three lines of evidence led us to think that CCA1 and LHY might drive circadian regulation of *CBF1*, *CBF2*, and *CBF3*. First, the protein levels for CCA1 and LHY peak in the early morning (ZT1–3) (26, 35, 36), just before the time that the transcript levels of *CBF1*, *CBF2*, and *CBF3* begin to increase. Second, the transcript levels for *PRR7* (32, 36) and *LIGHT-HARVESTING COMPLEX B (LHCB)* (24, 35, 37), both of which are induced by CCA1 and LHY, peak similarly to the *CBF* genes. Last, the promoter regions of *CBF1*, *CBF2*, and *CBF3* have several EE (AAAATATCT) (19) and CCA1-binding sites (CBS; AATCT) (35) (Fig. 2) that mediate binding of CCA1 and LHY (30, 35, 38) to target promoters. To determine whether CCA1 and LHY were involved in circadian regulation of *CBF1*, *CBF2*, and *CBF3*, we asked whether their expression was affected in plants carrying either the single *cca1-11* or *lhy-21* null mutations or the *cca1-11/lhy-21* double mutation. We found that the single mutations had dif-




Fig. 2. Binding of CCA1 at the *CBF1-3* locus. *cca1-1* and *cca1-1 CCA1p:CCA1-GFP* plants were grown at 22 °C under a 12-h photoperiod to the four-leaf stage. Tissue was fixed at ZT4, and ChIP was performed using anti-GFP antibody. Immunoprecipitated DNA was quantified by qRT-PCR using primers specific to regions within the *CBF1-3* locus (boxes A through O). The levels of immunoprecipitated DNA were normalized to the respective input DNA. Immunoprecipitation in *cca1-1 CCA1p:CCA1-GFP* plants (black bars) is relative to *cca1-1* plants (gray bars) set to a value of 1. Primer pairs directed to the 3' UTR of *TOC1* (*TOC1-3'UTR*), *ACTIN 7*, and *UBIQUITIN 10 (UBQ10)* were used as negative controls (31). Primers near the EE element in the *TOC1* promoter (*TOC1-EE*) were used as a positive control. Values represent the average of five independent biological experiments. Error bars indicate SEM. **P* < 0.05 using a paired, one-tailed *t* test. In the *CBF* locus diagram, the transcribed regions are indicated by white boxes, and the approximate positions of CBS [(A₂₋₄)TCT], and EE (AAAATATCT) motifs are indicated by gray circles and white squares, respectively.

fering effects on the three *CBF* genes, whereas cycling of *CBF1* was severely disrupted, *CBF2* and *CBF3* transcript levels cycled with approximately the same amplitudes as in the wild-type plants, although the peaks occurred about 2 h earlier than in the wild-type plants (Fig. S1). Period shortening of output genes such as *LHCB* has been observed previously in *cca1* and *lhy* mutant plants (39, 40) as well as in our experiments (Fig. S2). The *cca1-11/lhy-21* double mutations also had differing effects on the circadian regulation of the three *CBF* genes (Fig. 1); circadian regulation of *CBF1* and *CBF3* was essentially eliminated in the double-mutant plants, but *CBF2* transcript levels clearly continued to cycle, although the amplitude was diminished, and the period was greatly shortened. Period shortening of output genes such as *LHCB* also has been observed previously in *cca1-11/lhy-21* double-mutant plants (40, 41) and in our experiments (Fig. S3). From these results we concluded that circadian regulation of *CBF1* and *CBF3* is dependent on the action of either CCA1 or LHY and that circadian regulation of *CBF2* involves action of CCA1 and LHY but can be driven to a considerable degree by other unknown factors.

CCA1 and LHY may impart circadian regulation of *CBF1*, *CBF2*, and *CBF3* by binding to the EE and CBS motifs present in the *CBF* promoters and act as positive regulators stimulating transcription. To test this hypothesis, we conducted ChIP experiments to determine whether CCA1 binds directly to the promoter regions of *CBF1*, *CBF2*, and *CBF3*. To do so, we compared ChIP results obtained with plants carrying the *cca1-1* mutation and *cca1-1* plants that had been restored with a construct encoding the CCA1 protein tagged with GFP under the endogenous CCA1 promoter (31). Chromatin was isolated from plants harvested at ZT4, the point when *CBF* transcript levels begin to rise. In mock experiments where rabbit Ig was used for precipitation, no specific binding was detected for any of the subregions (A to O in Fig. S4) of the *CBF1-3* locus tested or for *TOC1*, *ACTIN 7*, or *UBIQUITIN 10* (Fig. S4). In contrast, test experiments indicated that specific binding of the CCA1-GFP protein occurred throughout most of the *CBF1-3* locus and also within the promoter region of *TOC1*, a positive control, but not in the promoters of *ACTIN 7* or *UBIQUITIN 10*, two negative controls (31) (Fig. 2). Significant CCA1-GFP binding occurred in the promoter regions of *CBF1*, *CBF2*, and *CBF3* (C, L, and G and I, respectively in Fig. 2). We also observed CCA1 associated with the coding region of *CBF1* (E in Fig. 2), perhaps because of the tandem connection of the *CBF* genes and consequently close proximity to several CCA1 binding sites located in the adjacent *CBF3* promoter (Fig. 2). Thus, although it is possible that there is CCA1 binding within the *CBF1* transcript region, the actual binding site could be downstream in the *CBF3* promoter. In sum, the results of our genetic and ChIP experiments support the model that circadian regulation of *CBF1*, *CBF2*, and *CBF3* involves action of CCA1 and LHY binding to the promoters of these genes and up-regulating their transcription during the morning hours.

Rhythmic Expression of CBF Regulon COR Genes and Freezing Tolerance Are Impaired in Plants Carrying the *cca1-11/lhy-21* Double Mutation. Harmer et al. (19) suggested that the circadian regulation of *CBF1*, *CBF2*, and *CBF3* could result in rhythmic expression of CBF regulon *COR* genes. To test this possibility, we examined the transcript levels for three CBF-inducible genes—*COR15A*, *COR47*, and *COR78*—in wild-type plants and in plants carrying the *cca1-11/lhy-21* double mutation. The results indicated that the transcript levels for all three *COR* genes oscillated with a period of about 24 h in wild-type plants, although for *COR15A* and *COR78* the amplitude of the second peak was much less than that of the first peak (Fig. 1). For all three genes, the first peak occurred at about ZT10, consistent with the transcript levels of *CBF1*, *CBF2*, and *CBF3* peaking just before this time, at about ZT8 (19, 20) (Fig. 1). Moreover, the oscillation in *COR* transcript levels was largely reduced in the *cca1-11/lhy-21* double-mutant plants (Fig. 1). These results were consistent

with the model that circadian-regulated expression of *CBF1*, *CBF2*, and *CBF3* imparts rhythmic expression of CBF-targeted *COR* genes at “basal” nonacclimating temperatures.

The decrease in expression of the CBF-targeted *COR* genes in the *cca1-11/lhy-21* double-mutant plants could result in a decrease in basal freezing tolerance. We tested this possibility by using the electrolyte leakage assay to compare the freezing tolerance of wild-type and *cca1-11/lhy-21* plants. The results indicated that the *cca1-11/lhy-21* double mutation reduced freezing tolerance by about 50%; although the temperature at which cell damage results in release of 50% of total electrolytes (EL₅₀) was about −4 °C in wild-type plants, it was about −2 °C in the *cca1-11/lhy-21* mutant plants (Fig. 3). Thus, the circadian clock is required for maximum basal freezing tolerance.

CCA1 and LHY Regulate Cold Induction of *CBF1*, *CBF2*, and *CBF3*.

Fowler et al. (22) reported that cold induction of *CBF1*, *CBF2*, and *CBF3* is gated by the circadian clock. Given our results indicating a role for CCA1 and LHY in the circadian regulation of *CBF1*, *CBF2*, and *CBF3*, we asked whether these transcription factors also had a role in the gating phenomenon. As previously reported (22, 34), cold-induction of *CBF1*, *CBF2*, and *CBF3* was much greater in the subjective day than in the subjective evening (Fig. 4). This cold induction was little affected by the single *cca1-11* and *lhy-21* mutations (Fig. S1) but was greatly reduced in the *cca1-11/lhy-21* double mutants, and the period of cycling was shortened (Fig. 4), as was the period of cycling for *LHCB* (Fig. S3). Thus,




Fig. 3. Effect of the *cca1-11/lhy-21* double mutation on plant freezing tolerance. *cca1-11/lhy-21* double-mutant and wild-type *Ws-2* (WS) plants were grown at 22 °C under a 12-h photoperiod for ~18 d and were tested directly for freezing tolerance (nonacclimated plants; Upper) or were transferred at ZT4 to 4 °C for 7 d under a 12-h photoperiod and then tested for freezing tolerance (cold-acclimated plants; Lower). Freezing tolerance was tested using the electrolyte leakage test. The results presented are average values from three independent experiments. Error bars indicate SEM.

CCA1 and LHY have a major role in the induction of *CBF1*, *CBF2*, and *CBF3* expression in response to low temperature.

Cold induction of CBF-Targeted COR Genes and Freezing Tolerance Are Impaired in Plants Carrying the *cca1-11/lhy-21* Double Mutation.

The finding that cold induction of *CBF1*, *CBF2*, and *CBF3* was impaired in plants carrying the *cca1-11/lhy-21* double mutation prompted us to determine whether the double mutation also impaired cold induction of CBF-targeted *COR* genes and the freezing tolerance of cold-acclimated plants. In wild-type plants, the degree to which *COR15A*, *COR47*, and *COR78* were induced by low temperature cycled, with peaks in the late day and troughs in the late subjective evening (Fig. 4), times that were consistent with the cycling of *CBF1*, *CBF2*, and *CBF3* cold induction (Fig. 4). In plants carrying the *cca1-11/lhy-21* double mutation, the peak following the night-to-day transition was little affected, but the subsequent peaks were greatly diminished, and the period of cycling was shortened (Fig. 4). In addition, the freezing tolerance of cold-acclimated plants carrying the *cca1-11/lhy-21* double mutation was about 1 °C less than that of cold-acclimated wild-type plants (Fig. 3). These results indicate that CCA1 and LHY are required for *Arabidopsis* plants to attain maximum levels of *COR* gene induction and freezing tolerance in response to low temperature.

Discussion

The CBF cold-response pathway is highly conserved among plants and has a major role in plant freezing tolerance (1–3, 9). Accordingly, there is considerable interest in understanding the mechanisms that control expression of this stress-response pathway. Here we establish that the CBF pathway is subject to positive regulation by the circadian clock components CCA1 and LHY. We show that these factors have roles in both circadian regulation and cold induction of the pathway and that they are required for plants to attain maximum freezing tolerance at both basal and cold-acclimating temperatures.

At basal growth temperature, the transcript levels for *CBF1*, *CBF2*, and *CBF3* oscillate, with peaks and troughs occurring at about ZT8 and ZT20, respectively (19–21) (Fig. 1). Our genetic and ChIP analyses indicate that this circadian regulation is caused by

the direct action of CCA1 and LHY binding at the *CBF1-3* locus—presumably at the EE, CBS, and related motifs—and induces transcription of the *CBF* genes. In the morning hours, when CCA1 and LHY protein levels peak (25, 35, 36), the transcript levels of *CBF1*, *CBF2*, and *CBF3* peak; in the evening hours, when CCA1 and LHY protein levels are low, the transcript levels for *CBF1*, *CBF2*, and *CBF3* are low (Fig. 5). The finding that circadian regulation of *CBF1* and *CBF3* is nearly eliminated in plants carrying the *cca1-11/lhy-21* double mutation indicates that no other regulatory proteins are sufficient to impart positive circadian regulation of these genes. In contrast, circadian regulation of *CBF2* continues in plants carrying the *cca1-11/lhy-21* double mutation, albeit with reduced amplitude and shortened periodicity. Thus, at least one additional regulatory protein appears to drive positive circadian regulation of *CBF2*. Prime candidates for this residual regulation are the four REVEILLE (RVE) proteins RVE1, RVE3, RVE4, and RVE8 (42). These Myb-like transcription factors fall into the CCA1 subfamily, bind to the EE motif, are circadian regulated, and, like CCA1 and LHY, have peak transcript levels at dawn (42).

Kidokoro et al. (21) reported that circadian regulation of *CBF1*, *CBF2*, and *CBF3* also involves negative regulation. These investigators found that PHYTOCHROME INTERACTING FACTOR 7 (PIF7) binds to a G-box element in the promoter of *CBF2* and that this element is required for down-regulation of the *CBF2* promoter during the subjective evening. In addition, they found that PIF7 physically interacts with TOC1 (21). Thus, circadian-controlled down-regulation of the *CBF* genes appears to involve action of a PIF7–TOC1 protein complex binding to G-box elements in their promoters (Fig. 5).

In addition to establishing a role for CCA1 and LHY in circadian regulation of *CBF1*, *CBF2*, and *CBF3*, our results indicate that CCA1 and LHY also act as positive regulators of *CBF* cold induction. This action is evidenced by the finding that cold induction of *CBF1*, *CBF2*, and *CBF3* is greatly impaired in plants carrying the *cca1-11/lhy-21* double mutation (Fig. 4). We propose that the gating of *CBF1*, *CBF2*, and *CBF3* cold induction results, in part, from positive synergistic interaction between cold-signaling and clock-output pathways, the former mediated by ICE1 and CAMTA3 and the latter by CCA1 and LHY (Fig. 5). If the




Fig. 4. Effects of the *cca1-11/lhy-21* double mutation on cold induction of *CBF1*, *CBF2*, *CBF3*, and CBF-targeted genes *COR15A*, *COR47*, and *COR78*. Wild-type *Ws-2* (WS) and *cca1-11/lhy-21* double-mutant plants were grown at 22 °C under a 12-h photoperiod to the four-leaf stage and then were transferred to constant light at ZT0 (subjective day and night are indicated by white and gray bars, respectively). Plants were transferred to cold temperature (4 °C) for 2 h, every 2 h (*CBF* genes) or for 4 h every 4 h (*COR* genes) at the start of constant-light conditions. Transcript levels for the indicated genes were determined by qRT-PCR. The x axis represents the time (ZT) when plants were transferred to cold temperature. Gene expression was normalized to UBQ10 for each sample. Gene expression is relative to one wild-type sample set to a value of 1 for each biological replicate. Values are averages from three independent biological experiments. Error bars indicate SEM. Primer pair sequences are given in Table S1.

1. Thomashow MF (1999) Plant cold acclimation: Freezing tolerance genes and regulatory mechanisms. *Annu Rev Plant Physiol Plant Mol Biol* 50:571–599.
2. Ruelland EVM, Zachowski A, Vaughn H (2009) Cold signaling and cold acclimation in plants. *Adv Bot Res* 49:36–54.
3. Chinnusamy V, Zhu J, Zhu JK (2007) Cold stress regulation of gene expression in plants. *Trends Plant Sci* 12:444–451.
4. Maruyama K, et al. (2004) Identification of cold-inducible downstream genes of the Arabidopsis DREB1A/CBF3 transcriptional factor using two microarray systems. *Plant J* 38:982–993.
5. Hannah MA, Heyer AG, Hinch DK (2005) A global survey of gene regulation during cold acclimation in *Arabidopsis thaliana*. *PLoS Genet* 1:e26.
6. Fowler S, Thomashow MF (2002) Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell* 14:1675–1690.
7. Chew YH, Halliday KJ (2010) A stress-free walk from Arabidopsis to crops. *Curr Opin Biotechnol* 22:1–6.
8. Hua J (2009) From freezing to scorching, transcriptional responses to temperature variations in plants. *Curr Opin Plant Biol* 12:568–573.
9. Thomashow MF (2010) Molecular basis of plant cold acclimation: Insights gained from studying the CBF cold response pathway. *Plant Physiol* 154:571–577.
10. Riechmann JL, Meyerowitz EM (1998) The AP2/EREBP family of plant transcription factors. *Biol Chem* 379:633–646.
11. Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF (2005) Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of Arabidopsis. *Plant J* 41:195–211.
12. Liu Q, et al. (1998) Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in Arabidopsis. *Plant Cell* 10:1391–1406.
13. Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF (1998) Arabidopsis CBF1 overexpression induces COR genes and enhances freezing tolerance. *Science* 280:104–106.
14. Gilmour SJ, Fowler SG, Thomashow MF (2004) Arabidopsis transcriptional activators CBF1, CBF2, and CBF3 have matching functional activities. *Plant Mol Biol* 54:767–781.
15. Chinnusamy V, et al. (2003) ICE1: A regulator of cold-induced transcriptome and freezing tolerance in Arabidopsis. *Genes Dev* 17:1043–1054.
16. Doherty CJ, Van Buskirk HA, Myers SJ, Thomashow MF (2009) Roles for Arabidopsis CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. *Plant Cell* 21:972–984.
17. Miura K, et al. (2007) SIZ1-mediated sumoylation of ICE1 controls CBF3/DREB1A expression and freezing tolerance in Arabidopsis. *Plant Cell* 19:1403–1414.
18. Knight H, Trewavas AJ, Knight MR (1996) Cold calcium signaling in Arabidopsis involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell* 8:489–503.
19. Harmer SL, et al. (2000) Orchestrated transcription of key pathways in Arabidopsis by the circadian clock. *Science* 290:2110–2113.
20. Bieniawska Z, et al. (2008) Disruption of the Arabidopsis circadian clock is responsible for extensive variation in the cold-responsive transcriptome. *Plant Physiol* 147:263–279.
21. Kidokoro S, et al. (2009) The phytochrome-interacting factor PIF7 negatively regulates DREB1 expression under circadian control in Arabidopsis. *Plant Physiol* 151:2046–2057.
22. Fowler SG, Cook D, Thomashow MF (2005) Low temperature induction of Arabidopsis CBF1, 2, and 3 is gated by the circadian clock. *Plant Physiol* 137:961–968.
23. Pruneda-Paz JL, Kay SA (2010) An expanding universe of circadian networks in higher plants. *Trends Plant Sci* 15:259–265.
24. Harmer SL (2009) The circadian system in higher plants. *Annu Rev Plant Biol* 60:357–377.
25. Wang ZY, Tobin EM (1998) Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. *Cell* 93:1207–1217.
26. Schaffer R, et al. (1998) The late elongated hypocotyl mutation of Arabidopsis disrupts circadian rhythms and the photoperiodic control of flowering. *Cell* 93:1219–1229.
27. Mizoguchi T, et al. (2002) LHY and CCA1 are partially redundant genes required to maintain circadian rhythms in Arabidopsis. *Dev Cell* 2:629–641.
28. Alabadi D, Yanovsky MJ, Más P, Harmer SL, Kay SA (2002) Critical role for CCA1 and LHY in maintaining circadian rhythmicity in Arabidopsis. *Curr Biol* 12:757–761.
29. Matsushika A, Makino S, Kojima M, Mizuno T (2000) Circadian waves of expression of the APRR1/TOC1 family of pseudo-response regulators in *Arabidopsis thaliana*: Insight into the plant circadian clock. *Plant Cell Physiol* 41:1002–1012.
30. Alabadi D, et al. (2001) Reciprocal regulation between TOC1 and LHY/CCA1 within the Arabidopsis circadian clock. *Science* 293:880–883.
31. Pruneda-Paz JL, Breton G, Para A, Kay SA (2009) A functional genomics approach reveals CHE as a component of the Arabidopsis circadian clock. *Science* 323:1481–1485.
32. Farré EM, Harmer SL, Harmon FG, Yanovsky MJ, Kay SA (2005) Overlapping and distinct roles of PRR7 and PRR9 in the Arabidopsis circadian clock. *Curr Biol* 15:47–54.
33. Nakamichi N, et al. (2010) PSEUDO-RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in the Arabidopsis circadian clock. *Plant Cell* 22:594–605.
34. Nakamichi N, et al. (2009) Transcript profiling of an Arabidopsis PSEUDO RESPONSE REGULATOR arrhythmic triple mutant reveals a role for the circadian clock in cold stress response. *Plant Cell Physiol* 50:447–462.
35. Wang ZY, et al. (1997) A Myb-related transcription factor is involved in the phytochrome regulation of an Arabidopsis Lhcb gene. *Plant Cell* 9:491–507.
36. Hamilton EE, Kay SA (2008) SnapShot: Circadian clock proteins. *Cell* 135:368–368e1.
37. Mockler TC, et al. (2007) The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern matching, and promoter analysis. *Cold Spring Harb Symp Quant Biol* 72:353–363.
38. Green RM, Tobin EM (1999) Loss of the circadian clock-associated protein 1 in Arabidopsis results in altered clock-regulated gene expression. *Proc Natl Acad Sci USA* 96:4176–4179.
39. Hall A, et al. (2003) The TIME FOR COFFEE gene maintains the amplitude and timing of Arabidopsis circadian clocks. *Plant Cell* 15:2719–2729.
40. Gould PD, et al. (2006) The molecular basis of temperature compensation in the Arabidopsis circadian clock. *Plant Cell* 18:1177–1187.
41. Locke JCW, et al. (2005) Extension of a genetic network model by iterative experimentation and mathematical analysis. *Mol Syst Biol*, 10.1038/msb4100018.
42. Gong W, et al. (2008) The development of protein microarrays and their applications in DNA-protein and protein-protein interaction analyses of Arabidopsis transcription factors. *Mol Plant* 1:27–41.
43. Fukushima A, et al. (2009) Impact of clock-associated Arabidopsis pseudo-response regulators in metabolic coordination. *Proc Natl Acad Sci USA* 106:7251–7256.
44. Espinoza C, et al. (2010) Interaction with diurnal and circadian regulation results in dynamic metabolic and transcriptional changes during cold acclimation in Arabidopsis. *PLoS ONE* 5:e14101.
45. Alonso JM, et al. (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301:653–657.